


JOURNAL OF BIOPHOTONICS

www.biophotonics-journal.org



Impairments of cerebral blood flow microcirculation in rats brought on by cardiac cessation and respiratory arrest

Gennadii Pivchenko, Igor Kozlov, Viktor Dremmin, Dmitry Stavtsev, Evgeniya Seryogina, Ksenia Kandurova, Valery Shupletsov, Konstantin Lapin, Alexander Alekseyev, Sergey Kuznetsov, Alexander Bykov, Andrey Dunaev, Igor Meglinski

WILEY-VCH

JOURNAL OF BIOPHOTONICS

Editor-in-Chief



Jürgen Popp

Institute of Physical Chemistry &
Abbe Center of Photonics (ACP),
Friedrich-Schiller-University Jena
and
Leibniz Institute of Photonic Technology
Jena, Germany
E-mail: juergen.popp@leibniz-ipht.de
(Photo: Sven Döring / Leibniz-IPHT)

Assistant Editor



Michael Schmitt

Institute of Physical Chemistry,
Friedrich-Schiller-University Jena,
Germany
E-mail: jbp@uni-jena.de

Editorial Office

Phone: +49-3641-948367
Fax: +49-3641-948302
E-mail: jbp@uni-jena.de

Journal Publishing Manager



Regina Hagen

Wiley-VCH GmbH
Berlin, Germany
E-mail: jbp@wiley.com

www.biophotonics-journal.org

Submission

Online manuscript submission: <https://mc.manuscriptcentral.com/jbio>

Journal of Biophotonics is the first international journal dedicated to publishing original articles and reviews from the exciting field of biophotonics, i.e. the development and application of photonic technologies in particular for (bio)medicine, but also life- and environmental sciences. The journal offers a platform where technology developers (physicists, chemists, engineers, etc.) communicate with endusers (in particular research clinicians) and where the clinical practitioner learns about the latest tools for the diagnosis and therapy of diseases. As such, the journal is highly interdisciplinary, publishing innovative research in the field of light interaction with biological material. The coverage extends from fundamental research to specific developments, while also including the latest applications or clinical trials/case reports.

Editorial Board

Peter E. Andersen

Technical University of Denmark, Roskilde, Denmark

Stefan Andersson-Engels

Lund Institute of Technology, Cork, Ireland

Stephen Boppart

Beckman Inst. for Advanced Science and Technology, Urbana, IL, USA

Arthur Chiou

National Yang-Ming University, Taipei, Taiwan

Kishan Dholakia

School of Physics & Astronomy, St Andrews, UK

Alberto Diaspro

University of Genoa, Italy

Max Diem

Northeastern University, Boston, MA, USA

Wolfgang Drexler

Medical University, Vienna, Austria

Paul French

Imperial College, London, UK

Martin Frenz

University of Bern, Switzerland

Min Gu

Swinburne Univ. of Technology, Hawthorn, Vic., Australia

Stefan Hell

Max-Planck-Institute for Biophysical Chemistry,
Göttingen, Germany

Elizabeth M. C. Hillman

Columbia University, New York, USA

Jiming Hu

Wuhan University, China

Zhiwei Huang

Singapore

Satoshi Kawata

LaSIE, Osaka University, Japan

Karsten König

Saarland University, Saarbrücken, Germany

Laura Marcu

University of California, Davis, CA, USA

Dennis L. Matthews

University of California, Sacramento, USA

Francesco S. Pavone

LENS - European Laboratory for Non-Linear
Spectroscopy, Florence, Italy

Paras Prasad

The State University of New York, Buffalo, NY, USA

Halina Rubinsztein-Dunlop

University of Queensland, Brisbane, Australia

David Sampson

The University of Western Australia, Crawley, Australia

Albert Stolow

Steaie Inst. for Molecular Sciences, Ottawa, ONT, Canada

Katarina Svanberg

Lund University Hospital, Sweden

Attila Tárnok

IMISE, University Leipzig, Germany

Valery V. Tuchin

Saratov State University, Saratov, Russia


Siva Umaphathy

Indian Institute of Science, Bangalore, India

Brian Wilson

Ontario Cancer Institute, Toronto, ONT, Canada

JOURNAL OF BIOPHOTONICS



Publishers

Wiley-VCH GmbH
Rotherstr. 21, 10245 Berlin, Germany
E-mail: physics@wiley-vch.de

Journal Customer Services: For ordering information, claims and any enquiry concerning your journal subscription please go to <https://wolsupport.wiley.com/s/contactsupport?tabset-a7d10=2> or contact your nearest office.

Americas: E-mail: cs-journals@wiley.com;

Phone: +1 877 762 2974.

Europe, Middle East and Africa:

E-mail: cs-journals@wiley.com; Phone: +44 (0) 1865 778315; 0800 1800 536 (Germany).

Asia Pacific: E-mail: cs-journals@wiley.com;

Phone: +65 6511 8000.

Japan: For Japanese speaking support,

E-mail: cs-japan@wiley.com;

Visit our Online Customer Help at

<https://wolsupport.wiley.com/s/contactsupport?tabset-a7d10=2>.

Copyright Permission:

Permission to reproduce material can be obtained via RightsLink® directly from the article abstract on Wiley Online Library (TOOLS → 'Request Permission' link). For other requests please contact: RIGHTS-and-LICENCES@wiley-vch.de.

Journal Publishing Manager: Regina Hagen

Phone: +49 (0) 30 47 03 13 21

E-mail: jbp@wiley.com

Typesetter: SPi-Global, India

Information for subscribers

JOURNAL OF BIOPHOTONICS (Online ISSN: 1864-06489) is published monthly.

FTE-based pricing, based on the overall demand and on the size and type of the subscribing institution (academic, government, hospital or corporate, applies for this Journal. Institutional subscription prices for 2022 can be found on the journal homepage at "Subscribe to this journal". Prices are exclusive of tax. Asia-Pacific GST, Canadian GST/HST and European VAT will be applied at the appropriate rates. For more information on current tax rates, please go to www.wileyonlinelibrary.com/tax-vat. The institutional price includes online access to the current and all online back files to January 1st 2018, where available. For other pricing options, including access information and terms and conditions, please visit www.wileyonlinelibrary.com/access.

Wiley's Corporate Citizenship initiative seeks to address the environmental, social, economic, and ethical challenges faced in our business and which are important to our diverse stakeholder groups. Since launching the initiative, we have focused on sharing our content with those in need, enhancing community philanthropy, reducing our carbon impact, creating global guidelines and best practices for paper use, establishing a vendor code of ethics, and engaging our colleagues and other stakeholders in our efforts. Follow our progress at www.wiley.com/go/citizenship.

Wiley is a founding member of the UN-backed HINARI, AGORA, and OARE initiatives. They are now collectively known as Research4Life, making online scientific content available free or at nominal cost to researchers in developing countries. Please visit Wiley's Content Access - Corporate Citizenship site: <http://www.wiley.com/WileyCDA/Section/id-390082.html>.

Instructions for Authors including details on manuscript preparation, submission instructions and all other information are available at www.biophotonics-journal.org.

Online manuscript submission: via ScholarOne Manuscripts at <https://mc.manuscriptcentral.com/jbio>.

Journal of Biophotonics employs a plagiarism detection system. By submitting your manuscript to this journal you accept that your manuscript may be screened for plagiarism against previously published works.

Open Access: Journal of Biophotonics accepts articles for Open Access publication. Please visit <https://authorservices.wiley.com/author-resources/Journal-Authors/open-access/hybrid-open-access.html> and select author guidelines for further information about Open Access.

View this journal at www.wileyonlinelibrary.com/journal/JBIO.

Disclaimer: The Publisher and Editors cannot be held responsible for errors or any consequences arising from the use of information contained in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher and Editors of the products advertised.


Copyright and Copying: Copyright © 2021 Wiley-VCH GmbH. All rights reserved. No part of this publication may be reproduced, stored or transmitted in any form or by any means without the prior permission in writing from the copyright holder. Authorization to copy items for internal and personal use is granted by the copyright holder for libraries and other users registered with their local Reproduction Rights Organisation (RRO), e.g. Copyright Clearance Center (CCC), 222 Rosewood Drive, Danvers, MA 01923, USA (www.copyright.com), provided the appropriate fee is paid directly to the RRO. This consent does not extend to other kinds of copying such as copying for general distribution, for advertising or promotional purposes, for republishing, for creating new collective works or for resale. Permissions for such reuse can be obtained using the RightsLink "Request Permissions" link on Wiley Online Library. Special requests should be addressed to: permissions@wiley.com

ISSN 1864-0648 (Online)

Outer Front Cover

Despite the fact that both acute cardiac cessation and respiratory arrest are the reasons of brain ischemia, its hemodynamics, oxygenation and structural pathology differs. This article represents blood re-distribution and neuronal tissue degeneration during the first decade of minutes after the breathing and heart beating violation. The results show that cardiac cessation is much more life-threatening statement that provokes blood circulation abnormalities and causes strong signs of cerebral ischemia in compare with the respiratory arrest.


Further details can be found in the article by G. Piavchenko, I. Kozlov, V. Dremin, D. Stavtsev, E. Seryogina, K. Kandurova, V. Shupletsov, K. Lapin, A. Alekseyev, S. Kuznetsov, A. Bykov, A. Dunaev, and I. Meglinski (e202100216).



Inside Cover

Pulmonary cancer is one of the most malignancies worldwide. We introduce a novel three-dimensional Convolutional Neural Network (3D-PulCNN) to classify three pulmonary subtypes based on the microscopic hyperspectral image because it can supply spatial information as well as spectral information. Combining the difference of spectral curves and texture feature between the three subtypes, we can classify LUAD, LUSC and SCLC more accurately in cytodiagnosis.

Further details can be found in the article by Qing Zhang | Yan Wang | Song Qiu | Jiangang Chen | Li Sun and Qingli Li (e202100142).




RESEARCH ARTICLES

e202100216

Impairments of cerebral blood flow microcirculation in rats brought on by cardiac cessation and respiratory arrest

Gennadii Piavchenko*, Igor Kozlov, Viktor Dremmin, Dmitry Stavtsev, Evgeniya Seryogina, Ksenia Kandurova, Valery Shupletsov, Konstantin Lapin, Alexander Alekseyev, Sergey Kuznetsov, Alexander Bykov, Andrey Dunaev, Igor Meglinski*




After an impair of blood and respiration functions there are significantly different changes in the structure and function of brain tissue that we present in this article.

e202100142

3D-PulCNN: Pulmonary cancer classification from hyperspectral images using convolution combination unit based CNN




Qing Zhang, Yan Wang, Song Qiu, Jiangang Chen, Li Sun, Qingli Li*



Pulmonary cancer is one of the most common malignancies worldwide. Existing algorithms for its classification are mostly based on color images, and the improvement of accuracy is quite challenging. This article proposes a new framework named as three-dimensional convolutional neural network (3D-PulCNN) for classifying pulmonary cancer based on microscopic hyperspectral image. Also, 3D-UNet is leveraged to segment cancer cells, and their morphological characteristics are calculated to help with prognosis assessment.

RESEARCH ARTICLE

Impairments of cerebral blood flow microcirculation in rats brought on by cardiac cessation and respiratory arrest

Gennadii Piavchenko^{1,2,3*}  | Igor Kozlov⁴ | Viktor Dremine^{3,4,8}  |
 Dmitry Stavtsev^{4,6} | Evgeniya Seryogina³ | Ksenia Kandurova⁴ |
 Valery Shupletsov⁴ | Konstantin Lapin² | Alexander Alekseyev³ |
 Sergey Kuznetsov¹ | Alexander Bykov⁵ | Andrey Dunaev^{3,4} | Igor Meglinski^{1,2,5,7,8*} 

¹Department of Histology, Cytology and Embryology, I.M. Sechenov First Moscow State Medical University, Moscow, Russia

²V.A. Negovsky Scientific Research Institute of General Reanimatology, Federal Research and Clinical Centre of Intensive Care Medicine and Rehabilitology, Moscow, Russia

³Cell Physiology and Pathology Laboratory, Orel State University, Orel, Russia

⁴R&D Center of Biomedical Photonics, Orel State University, Orel, Russia

⁵Opto-Electronics and Measurement Techniques, Faculty of Information and Electrical Engineering, University of Oulu, Oulu, Finland

⁶Institute of Biomedical Systems, National Research University of Electronic Technology (MIET), Zelenograd, Russia

⁷Immanuel Kant Baltic Federal University, Kaliningrad, Russia

⁸College of Engineering and Physical Sciences, Aston University, Birmingham, UK

***Correspondence**

Gennadii Piavchenko and Igor Meglinski, 8/2 Trubetskaya str., Moscow, 119991, Russia.


Email: gennadii.piavchenko@staff.sechenov.ru and i.meglinski@aston.ac.uk

Funding information

H2020 Marie Skłodowska-Curie Actions, Grant/Award Numbers: 839888, 863214; Ministry of Science and Higher Education

Abstract

The impairments of cerebral blood flow microcirculation brought on by cardiac and respiratory arrest were assessed with multi-modal diagnostic facilities, utilising laser speckle contrast imaging, fluorescence spectroscopy and diffuse reflectance spectroscopy. The results of laser speckle contrast imaging show a notable reduction of cerebral blood flow in small and medium size vessels during a few minutes of respiratory arrest, while the same effect was observed in large sinuses and their branches during the circulatory cessation. Concurrently, the redox ratio assessed with fluorescence spectroscopy indicates progressing hypoxia, NADH accumulation and increase of FAD consumption. The results of diffuse reflectance spectra measurements display a more rapid grow of the perfusion of deoxygenated blood in case of circulatory impairment. In addition, consequent histopathological analysis performed by using new tissue staining procedure developed in-house. It shows notably higher reduction of size of the neurons due to their wrinkling within brain tissues influenced by circulation impair. Whereas, the brain tissues altered with the respiratory arrest demonstrate focal perivascular oedema and mild hypoxic changes of neuronal morphology. Thus, the study suggests that consequences of a cessation of cerebral blood flow become more dramatic and dangerous compare to respiratory arrest.



This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Journal of Biophotonics* published by Wiley-VCH GmbH.

of the Russian Federation, Grant/Award Number: 075-15-2020-926; Academy of Finland, Grant/Award Number: 326204

KEYWORDS

brain imaging, cardiac cessation, cerebral blood flow, laser speckle contrast, microcirculation, respiratory arrest

1 | INTRODUCTION

Nowadays, prediction of early damage and microstructural reorganisation of cerebral tissues composition is crucial for numerous acute states, and especially those associated with a risk of brain death. The clinical death is typically caused by cessation of blood circulation and/or respiratory arrest [1]. In current study, we combined use of modern photonics-based modalities, such as laser speckle contrast imaging (LSCI), fluorescence spectroscopy (FS) and diffuse reflectance spectroscopy (DRS), the impairments of cerebral blood flow microcirculation brought by cardiac cessation and respiratory arrest are comparatively investigated.

LSCI is well-known imaging modality [2], and used extensively in a number of blood microcirculation studies, including those of tumour angiogenesis [3], cutaneous vessels reaction to allergens [4, 5], skin complication of diabetes [6], respiratory-related blood flow oscillations [7], and visualisation of cerebral vascular network [8]. It has been also demonstrated that with a combined application of LSCI and intravital fluorescence imaging the transcranial visualisation of brain's vascular bed can be performed in terms of blood flow within single arteries and veins [9–11].

DRS is a promising inexpensive noninvasive method that is able to show real-time dynamics of acute cerebral pathologies and to assess cerebral haemodynamics [12]. When FS and DRS are successfully utilised for monitoring metabolic and morphological changes in biological tissues in vivo [13].

In addition to the optical methods, histopathology is widely used for quantitative assessment of brain damage and direct visualisation of structural malformations associated with diseases and brain death. Histopathological analysis, used for the pathological changes within brain tissues, includes both routine staining with hemotoxylin and eosin (HE) as well as own new method of staining. In case of classical approach of HE staining, we are able to see only the most expressed pathology, making conclusion questionable. With new method we detect strong neuronal ischemia and vascular disorders as well as two types of neurons stained with cresyl violet and silver impregnation. The neuronal cells vary in structural and tinctorial properties and indicate various types of pathologies.

Thus, LSCI and DRS are used for evaluation of impairments of cerebral blood flow and blood microcirculation brought on by acute hypoxia provoked by respiration arrest and cardiac cessation. The associated variations in metabolic activities in cerebral cortex are assessed quantitatively by FS. In addition, the histopathological analysis is used to detect a hypoxic impact on the cerebral cortex in terms of structural morphological changes in postmortem brain tissues.

2 | MATERIALS AND METHODS

The experimental system combining LSCI, DRS and FS imaging/diagnostic modalities is presented in Figure 1.

The laser source (10 mW, 785 nm operating wavelength, Thorlabs, Inc., USA) illuminates the area of interest through a diffuser. The back-scattered light is collected via the lens (MVL25TM23, Thorlabs, Inc., USA), and the laser speckles analysis is performed by high-resolution CMOS camera (DCC3260M, 1936 × 1216 pixels and 5.86-pixel size, Thorlabs, Inc., USA). To eliminate single scattering, two crossed polarizers are placed in front of camera objective and laser module (see Figure 1). The speckle contrast is counted for 40 Hz frame rate and 20 ms expose time, as [14]:

$$C = \frac{\sigma}{\langle I \rangle_k}, \quad (1)$$

where σ is the standard deviation for the intensity of the back scattered light, $\langle I \rangle_k$ is the mean intensity value, $\langle \dots \rangle$ denotes averaging by k values of the sequential stacks of images ($k = 40$). The spatial averaging of raw speckle image obtained with CMOS camera is performed utilising 5×5 pixels sliding window [10]. This approach aims to improve the signal-to-noise ratio and image quality. Speckle perfusion index (SFI) is defined as [15]:

$$SFI = \frac{1}{2TC^2}, \quad (2)$$

where T is the exposure time of camera.

Every frame of stack is normalised by frame with the lowest mean of SFI. Thus, relative SFI is calculated as:




FIGURE 1 Experimental multi-modal imaging system developed in-house (A): 1—CMOS camera, 2—camera lens, 3—785 nm laser, 4, 5—polarisers, 6—diffuser, 7—spectrometer, 8—filter holder, 9—365 nm LED, 10—450 nm laser diode, 11—halogen lamp, 12—needle probe; close-up look of the needle probe. Example of raw speckle image from rat brain surface (B), “A” denotes “anterior” and “P” denotes “posterior” anatomical directions; registered fluorescence (C) and diffuse reflectance spectra (D)

$$\text{relative SFI} = 100 \times \frac{\text{SFI}(t)}{\text{baseline SFI}}, \quad (3)$$

where *baseline SFI* is the averaged map by first 30 frames.


LSCI measurements are recorded at the same time with FS and DRS measurements [13]. The FS mode includes 365 nm LED and 450 nm laser diode to excite the fluorescence of NADH and FAD coenzymes [16, 17]. Optical filters are used with cut off wavelengths of 400 and 490 nm (Thorlabs Inc., USA), respectively, to attenuate the backscattered light. DRS mode includes 360 to 2400 nm tungsten halogen light source HL-2000-FHSA (Ocean Insight, USA). The FS and DRS spectra are recorded by “FLAME” spectrometer (Ocean Insight, USA) and analysed in the range of 400 to 900 nm. The custom application is developed in MATLAB (MathWorks, Inc., USA) environment for data recording, saving and controlling the setup.

The computational analysis of spatial localisation of the LSCI and FS detected signals [18, 19] at 785 nm shows that the effective penetration depth in rodents tissues lies in a range $\sim 0.5\text{--}0.7$ mm). Since the thickness of


skin and skull in rodents is about ~ 0.5 mm, the young rats have been selected for noninvasive transcranial cerebral imaging.

To deliver the optical radiation into the cortex, a custom fibre-optic probe is applied (see Figure 1). The probe has 1 mm diameter and 20° bevel to ensure a good contact as well as not to damage brain tissues significantly. The probe contains 10 optical fibres (0.22 NA). The central one (200 μm in diameter) the light from the tissue surface to the spectrometer [17], whereas the surrounding fibres (100 μm in diameter, three fibres per light source). To avoid uncertainties in spectral measurements associated with the local suppression of superficial soft tissues [20, 21] the probe is placed close to the skin surface without touching it. The FS and DRS spectra are recorded during 1.5 and 0.5 s, respectively. To avoid a photobleaching effect, fluorescence excitation sources have been turned off between the measurements. The examples of registered raw speckle image, FS and DRS are shown in Figure 1B–D. Both 2 weeks old and 2 months old male Wistar rats were used in the experiments. All animals were kept in controlled environmental conditions (20–26°C of temperature, 50–60% of humidity, 12 h day-night cycle and 10 times air exchange

Respiratory arrest model




Non-depolarizing muscle relaxant (cisatracurium) in a lethal dose



Immediately causes a spasm of the respiratory muscles and respiratory arrest

Cardiac cessation model

Lidocaine solution (2%) in a dose of 1 ml



Immediately causes a cardiac arrest




FIGURE 2 Schematic presentation of respiratory and cardiac cessation. After placement in the stereotaxic apparatus the animal was administrated with drugs that caused immediate respiratory arrest (propofol followed by cisatracurium) or acute cardiac cessation (lidocaine solution)




FIGURE 3 Blood flow dynamics in time. The upper row corresponds to respiration impair (A, B, C), and the lower one to cardiac cessation (D, E, F). Notable changes appear from 1st minute of the recording (B, E). Video for blood flow dynamics is available (Rat brain. Cardiac cessation and respiratory arrest, 191.4 Mb, .avi [URL: <https://doi.org/10.5281/zenodo.4739049>])

per hour) in accordance with the principles of Good Laboratory Practice [22].

The animals were anaesthetized by an intramuscular injection of Zoletil drug (dose—25 mg/kg, Vibrac, France) and catheterized into right external jugular vein with PM-60 polymer catheter (SciCat, Russia, $d_{\text{ext}} = 0.8$ mm, $d_{\text{int}} = 0.4$ mm). The jugular veins catheterization approach is anatomically easily accessible with a minimal influence on the circulatory system, and is also accompanied by a lower risk of blood loss. The injection of drug into the jugular vein in comparison with the tail vein provides better compatibility [23]. After the anaesthesia, the animal is placed on a heating table to control body temperature and is fixed in a 3D-printed animal body holder, developed in-house, to avoid movement artefacts [24]. All the manipulations with the animals are

approved by the ethical committee of the Orel State University named after I.S. Turgenev (protocol No. 10, 16 October 2018).

The experimental models of respiratory and cardiac arrest are performed for the corresponding animals groups by the injection of drugs combination via catheter (Figure 2). We have checked the heart rate (to confirm the cardiac cessation) with an electrode detecting the heart beats during the experiment. We also counted the respiratory rate (number of diaphragm contractions) by a visual registration of breathing acts. As far as cisatracurium immediately causes a respiratory muscles spasm, any last diaphragm contraction has been considered as a breathing stop in the animal.

After LSCI, FS, DRS imaging and spectra recording, brains of all the animals were fixed in buffered formalin. Paraffin embedded sections of 5 μm thickness are prepared

FIGURE 4 The LSCI images observed during respiratory arrest (A) and cardiac cessation (B) arrest with the selected areas of particular groups of vessels: 1—veins of medium calibre, 2—vessels of microcirculation, 3—venous sinus. Corresponded temporal variations of relative SFI (C, D, E) observed during respiratory (blue) and cardiac (red) arrest in the selected areas: 1, 2, 3. Anatomical direction signs are same as Figure 1B





FIGURE 5 The results of FS and DRS measurements: (A) redox ratios of fluorescence intensity (maxima associated with NADH and FAD); (B) DRS ratio obtained at 540 and 805 nm during respiration arrest (C) and cardiac cessation (D)



with Microm HM 450 microtome (Microm International GmbH, Germany) and stained with HE by standard protocols [25] and with our own patented Nissl staining and silver impregnation combined method [26]. This new approach allows to see both silver impregnated and cresyl violet stained neurons with different structural and functional properties. The prepared slides were studied with

Leica DM4 M Microscope (Leica Microsystems GmbH, Germany) focusing on finding the possible ischemia and vascular pathology changes in the motor cortex area, as this region mostly contributes in the motor behaviour regulation (functioning as motor-related areas of cortex [27]). Paxinos and Watson Rat Brain Stereotaxic Atlas [28] is used as a reference guideline.




FIGURE 6 The results of histopathological analysis of coronal brain tissue sections in vitro: (A, C) rat motor cortex stained with haematoxylin and eosin; (B, D) rat motor cortex stained with method developed in-house utilising cresyl violet and silver impregnation combination. The circles highlight vessels with perivascular oedema (1), cresyl violet stained neurons with nucleoli (2), totally impregnated neurons (3), necrobiotic cells (4), ghost-like cells (5)

3 | RESULTS AND DISCUSSION

The impairments of cerebral blood flow microcirculation in cardiac cessation and respiratory arrest are presented in Figure 3. Anatomical direction signs are same as Figure 1B. The cerebral blood flow microcirculation measured with LSCI at normal conditions are seen in Figure 3A,D. Figure 3B,C, E show the map of cerebral blood flow microcirculation after 1 min of respiratory arrest and cardiac cessation, respectively, whereas Figure 3C,F represent same mappings after 2 min passed.

Temporal variations of relative SFI observed for the areas which correspond to the large (venous sinus, $\sim 100 \mu\text{m}$ in diameter), medium (veins of medium calibre, $\sim 30\text{--}40 \mu\text{m}$ in diameter) and small (vessels of microcirculation, $\sim 2\text{--}10 \mu\text{m}$ in diameter) cerebral blood vessels (Figure 4A,B) are presented in Figure 4C–E. As one can see, the relative SFI becomes steady for the first 30 s of measurements for both models. While, for the respiration arrest it takes 1 min (0.5–1.5 min) to get an extensive decrease of the relative SFI while in case of cardiac cessation it occurs in few seconds (see Figure 4). Further, a monotonic decrease of the relative SFI is observed with a flop between 6 and 8 min of observation. The obtained results suggest that for the respiration arrest the blood flow becomes most influenced first in medium and large size vessels (see Figure 4C and E, respectively), with the following monotonic decrease of flow. Whereas in small vessels the intensity of blood microcirculation decreasing after first minute is preserved both in case of respiratory arrest and cardiac cessation (see Figure 4D). The blood flow in the venous sinus is dropped significantly and much faster in case of cardiac cessation (see Figure 4E).

The results of FS and DRS measurements analysed with Origin Pro (OriginLab Corp., USA) and MATLAB, are presented in Figure 5. In the FS measurements, the maxima of emission intensity at 490 to 510 nm (for the excitation wavelength of 365 nm) and 510 to 530 nm (for 450 nm), respectively, are caused by the dominant contribution of NADH (IF_{NADH}) and FAD (IF_{FAD}). The redox ratio (RR) [29] defined as [30, 31]:

$$RR = \frac{IF_{\text{NADH}}}{IF_{\text{FAD}}} \quad (4)$$

is used to evaluate a metabolic changes of brain tissues associated with the impairments of cerebral blood flow microcirculation brought by cardiac cessation and respiratory arrest.

The RR demonstrates a progressive increase in hypoxia, an accumulation of NADH and increased FAD consumption. Trends identification for the full duration of the experiment is carried out using an exponential fitting with a R-square evaluation. These results are well agreed with the results of an alternative study [32, 33]. The changes of RR are more expressed for the case of respiration arrest, and showing less influence in the acute cerebral cortex ischemia due to cardiac cessation. The acute intensity changes are observed for the first 5 min after the heart arrest. The increase of RR is caused by growth of NADH-associated fluorescence and decrease of FAD. The obtained result suggests that blood circulation fail leads to a more acute progression of ischemia and to following metabolic derangements in brain cortex.

The DRS measurements at the range of wavelengths associated with oxy- (540 nm) and deoxy- (560 nm)

absorption peaks of haemoglobin, as well as at the isobestic point of 805 nm of oxy- and doxy-absorption provides an extra opportunity to assess the changes in blood content [34]. The results of DRS measurements show relatively steady dynamics of the reflectance ratio between maximum and minimum values, that do not exceed 5% (see Figure 5B). Both cases demonstrate reflectance changes (see Figure 5C,D) that are associated with the decrements of oxyhemoglobin within the sampling volume. These changes have no significant impact on the results of FS which are influenced by metabolic changes due to hypoxia [35, 36].

In addition to the results of LSCI, FS and DRS measurements mentioned above, the histopathology analysis of brain tissues is presented in Figure 6. In the brain cortex sections of the first respiration arrest group, a focal perivascular oedema and mild hypoxic changes of neuronal morphology are clearly seen (see Figure 6A). In the case of cardiac cessation, the brain cortex contains a significant number of hyperchromic wrinkled neurons, glial nodules at the site of necrotic neurons, as well as pericellular oedema in all tissue sections (see Figure 6C).

For specimens stained with the new approach developed in-house, in both cases the pathological changes in nervous tissue morphology are observed (see Figure 6B,D). The results of histopathology analysis for the group of animals with cardiac cessation also show a more considerable size reduction of impregnated and cresyl violet stained neurons due to their wrinkling. There are also initial stages of anucleated pale “ghost-like” neurons forming, a less distinct neuropil structure, a decreased basophilia of cellular cytoplasm (due to the reduction of activity in synthetic apparatus, or chromatophilic substance), a less smoothed outline of the neurons as well as dark impregnated neurons, (see Figure 6B,D). The new staining approach allows to compare the changes in neuronal morphology provoked by impairments of cerebral blood flow microcirculation brought by cardiac cessation and respiratory arrest.

4 | SUMMARY AND CONCLUSIONS

In acute respiratory arrest there is a gradual increase of hypoxia as well as a slowdown of haemodynamics. For acute circulatory impair we observe an immediate cease of oxygenated blood supply to the brain cortex. We conclude that acute cardiovascular impair (e.g., sudden cardiac arrest) is more life-threatening and leads to a faster increase of neuronal metabolism deficiency in comparison with acute respiratory impair. After long time hypoxia, the biochemical processes shift to the low energy mode and provide a resistance to hypoxia until a state of full anoxia

is achieved. The obtained results are well agreed with known literature data. Changes of FS intensities, which lead to a RR increase, might be noticed after 5 min, while the trend for DRS ratios remains almost the same, especially in the case of respiration impair. Blood circulation in vessels of cerebral cortex stops in case of respiratory arrest after about 2 min while in heart arrest it happens after about 1 min. Changes in the brain tissue oxygen saturation for heart arrest become evident during the first 5 min, but for acute breathing fail there are almost no changes. Heart arrest manifests by a more acute hypoxic state in cerebral cortex comparing to what is happening when breathing is impaired. So, acute cardiovascular impair is more life-threatening. It causes a rapid increase in neuronal metabolic deficiency comparing to acute respiratory impair. This may serve to personalise the management of patients after acute circulatory and respiratory arrest for prognosis and rehabilitation. The utilised multimodal photonics-based approach of brain tissue functional monitoring may allow to bring a complex assessment with laser speckle and spectroscopy methods for evaluation of brain metabolism changes in neurosurgical patients.

ACKNOWLEDGMENTS

The authors disclose receipt of financial support for the research, authorship, and/or publication of this article as following: Gennadii Pivachenko and Sergey Kuznetsov acknowledge the support from the Ministry of Science and Higher Education of the Russian Federation within the framework of State support for the creation and development of World-Class research Centres “Digital Biodesign and Personalized Healthcare” No 075-15-2020-926. Viktor Dremin kindly acknowledges personal support from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 839888. Andrey Dunaev and Igor Meglinski acknowledge funding from the Academy of Finland (grant No. 326204). This study has been also partially supported by the European Union’s Horizon 2020 research and innovation programme under grant agreement No. 863214—NEUROPA project.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Gennadii Pivachenko  <https://orcid.org/0000-0001-7782-3468>

Viktor Dremin  <https://orcid.org/0000-0001-6974-3505>

Igor Meglinski  <https://orcid.org/0000-0002-7613-8191>

REFERENCES

- [1] A. Walker, *Cerebral death*, 3rd ed., Urban & Schwarzenberg, Baltimore, MD **1985**.
- [2] W. Heeman, W. Steenbergen, G. van Dam, E. C. Boerma, *J Biomed Opt.* **2019**, *24*(8), 080901.
- [3] V. Kalchenko, N. Madar-Balakirski, I. Meglinski, A. Harmelin, *J. Biophotonics* **2011**, *4*(9), 645.
- [4] V. Kalchenko, Y. Kuznetsov, D. Preise, I. Meglinski, A. Harmelin, *J Biomed Opt.* **2014**, *19*(6), 060502.
- [5] V. Kalchenko, I. Meglinski, A. Sdobnov, Y. Kuznetsov, A. Harmelin, *J Biomed Opt* **2019**, *24*(6), 060501.
- [6] E. Zharkikh, V. Dremin, E. Zherebtsov, A. Dunaev, I. Meglinski, *J. Biophoton* **2020**, *13*(10), e202000203.
- [7] I. Mizeva, E. Zharkikh, V. Dremin, E. Zherebtsov, I. Makovik, E. Potapova, A. Dunaev, *Microvasc. Res.* **2018**, *120*, 13.
- [8] V. Kalchenko, D. Israeli, Y. Kuznetsov, A. Harmelin, *Sci. Rep.* **2014**, *4*(1), 5839.
- [9] V. Kalchenko, D. Israeli, Y. Kuznetsov, I. Meglinski, A. Harmelin, *J. Biophotonics* **2015**, *8*(11–12), 897.
- [10] V. Kalchenko, A. Sdobnov, I. Meglinski, Y. Kuznetsov, G. Molodij, A. Harmelin, *Photonics* **2019**, *6*(3), 80.
- [11] G. Molodij, A. Sdobnov, Y. Kuznetsov, A. Harmelin, I. Meglinski, V. Kalchenko, *Phys. Med. Biol.* **2020**, *65*(7), 075007.
- [12] R. M. Forti, M. Katsurayama, J. Menko, L. Valler, A. Quiroga, A. L. E. Falcão, L. M. Li, R. C. Mesquita, *Front. Med.* **2020**, *7*, 147.
- [13] V. Dremin, E. Potapova, E. Zherebtsov, K. Kandaurova, V. Shupletsov, A. Alekseyev, A. Mamoshin, A. Dunaev, *Sci. Rep.* **2020**, *10*, 14200.
- [14] J. D. Briers, S. Webster, *J Biomed Opt.* **1996**, *1*(2), 174.
- [15] C. Crouzet, R. H. Wilson, A. Bazrafkan, M. H. Farahabadi, D. Lee, J. Alcocer, B. J. Tromberg, B. Choi, Y. Akbari, *Biomed. Opt. Express* **2016**, *7*(11), 4660.
- [16] A. C. Croce, G. Bottiroli, *Eur J Histochem* **2014**, *58*(4), 2461.
- [17] A. V. Dunaev, V. V. Dremin, E. A. Zherebtsov, I. E. Rafailov, K. S. Litvinova, S. G. Palmer, N. A. Stewart, S. G. Sokolovski, E. U. Rafailov, *Med. Eng. Phys.* **2015**, *37*(6), 574.
- [18] Y. L. Kuznetsov, V. V. Kalchenko, N. G. Astafeva, I. V. Meglinski, *Quant. Electron* **2014**, *44*(8), 713.
- [19] V. Dremin, E. Zherebtsov, A. Bykov, A. Popov, A. Doronin, I. Meglinski, *Appl. Optics* **2019**, *58*(34), 9398.
- [20] A. Popov, A. Bykov, I. Meglinski, *J Biomed Opt* **2017**, *22*(11), 110504.
- [21] I. Mizeeva, E. Potapova, V. Dremin, E. Zherebtsov, M. Mezentsev, V. Shuleptsov, A. Dunaev, *Clin. Hemorheol. Microcirc.* **2019**, *72*(3), 259.
- [22] OECD Series on principles of good laboratory practice and compliance monitoring. <https://www.oecd.org/officialdocuments/publicdisplaydocumentpdf/?cote=env/mc/chem%2898%2917&doclanguage=en>.
- [23] K. V. Thirivikraman, R. L. Huot, P. M. Plotsky, *Brain Res. Brain Res. Protoc* **2002**, *10*(2), 84.
- [24] I. M. Gryadunov, G. A. Piavchenko, I. O. Kozlov, E. S. Seryogina, S. L. Kuznetsov, RU patent 200025U1 (**2020**).
- [25] G. A. P'yavchenko, L. I. Shmarkova, V. I. Nozdrin, *Neurosci. Behav. Physiol* **2016**, *46*(3), 270.
- [26] G. A. Piavchenko, D. Pranab, N. S. Novikova, V. I. Nozdrin, RU patent 2666256C2 (**2018**).
- [27] X. Li, M. A. Krol, S. Jahani, D. A. Boas, H. Tager-Flusberg, M. A. Yücel, *Sci. Rep* **2020**, *10*(1), 10965.
- [28] G. Paxinos, C. Watson, *The Rat Brain in Stereotaxic Coordinates: Hard Cover Edition*, Elsevier, San Diego, CA **2006**.
- [29] B. Chance, B. Schoener, R. Oshino, F. Itshak, Y. Nakase, *J. Biol. Chem* **1979**, *254*(11), 4764.
- [30] J. H. Ostrander, C. M. McMahon, S. Lem, S. R. Millon, J. Q. Brown, V. L. Seewaldt, N. Ramanujam, *Cancer Res.* **2010**, *70*(11), 4759.
- [31] K. Staniszewski, S. H. Audi, R. Sepehr, E. R. Jacobs, M. Ranji, *Ann. Biomed. Eng.* **2013**, *41*(4), 827.
- [32] V. Shupletsov, K. Kandaurova, E. Seryogina, G. Piavchenko, V. Dremin, A. Mamoshin, A. Dunaev, *Proc. SPIE* **2019**, *11140*, 1114001-212.
- [33] V. Dremin, E. Potapova, A. Mamoshin, A. Dunaev, E. Rafailov, *Laser Phys. Lett* **2020**, *17*(11), 115605.
- [34] B. Wilson, S. Jacques, *IEEE J Quant Electron* **1990**, *26*(12), 2186.
- [35] A. I. Zherebtsova, V. Dremin, I. N. Makovik, E. A. Zherebtsov, A. V. Dunaev, A. Goltsov, S. G. Sokolovski, E. U. Rafailov, *Front. Physiol.* **2019**, *10*, 416.
- [36] V. V. Dremin, E. A. Zherebtsov, V. V. Sidorov, A. Krupatkin, I. N. Makovik, A. I. Zherebtsova, E. V. Zharkikh, E. V. Potapova, A. V. Dunaev, A. A. Doronin, A. V. Bykov, I. E. Rafailov, K. S. Litvinova, S. G. Sokolovski, E. U. Rafailov, *J Biomed Opt.* **2017**, *22*(8), 1.

How to cite this article: G. Piavchenko, I. Kozlov, V. Dremin, D. Stavtsev, E. Seryogina, K. Kandaurova, V. Shupletsov, K. Lapin, A. Alekseyev, S. Kuznetsov, A. Bykov, A. Dunaev, I. Meglinski, *J. Biophotonics* **2021**, *14*(12), e202100216. <https://doi.org/10.1002/jbio.202100216>